



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Cycle threshold values are inversely associated with poorer outcomes in hospitalised patients with Covid-19**

**Citation for published version:**

Wright, J, Achana, F, Diwakar, L, Semple, MG, Carroll, WD, Baillie, K, Thompson, C, Alcock, A & Kemp, TS 2021, 'Cycle threshold values are inversely associated with poorer outcomes in hospitalised patients with Covid-19: a prospective, observational cohort study conducted at a UK tertiary hospital', *International Journal of Infectious Diseases*. <https://doi.org/10.1016/j.ijid.2021.08.022>

**Digital Object Identifier (DOI):**

[10.1016/j.ijid.2021.08.022](https://doi.org/10.1016/j.ijid.2021.08.022)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Version created as part of publication process; publisher's layout; not normally made publicly available

**Published In:**

International Journal of Infectious Diseases

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



Cycle threshold values are inversely associated with poorer outcomes in hospitalised patients with Covid-19: a prospective, observational cohort study conducted at a UK tertiary hospital.

Jenny Wright , Felix Achana , Lavanya Diwakar ,  
Malcolm G Semple , Will D Carroll , Kenneth Baillie ,  
Christopher Thompson , Alice Alcock , Timothy S Kemp

PII: S1201-9712(21)00657-3  
DOI: <https://doi.org/10.1016/j.ijid.2021.08.022>  
Reference: IJID 5634



To appear in: *International Journal of Infectious Diseases*

Received date: 18 June 2021  
Revised date: 11 August 2021  
Accepted date: 12 August 2021

Please cite this article as: Jenny Wright , Felix Achana , Lavanya Diwakar , Malcolm G Semple , Will D Carroll , Kenneth Baillie , Christopher Thompson , Alice Alcock , Timothy S Kemp , Cycle threshold values are inversely associated with poorer outcomes in hospitalised patients with Covid-19: a prospective, observational cohort study conducted at a UK tertiary hospital., *International Journal of Infectious Diseases* (2021), doi: <https://doi.org/10.1016/j.ijid.2021.08.022>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd on behalf of International Society for Infectious Diseases.  
This is an open access article under the CC BY-NC-ND license  
(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

**Title:**

Cycle threshold values are inversely associated with poorer outcomes in hospitalised patients with Covid-19: a prospective, observational cohort study conducted at a UK tertiary hospital.

**Authors:**

Jenny Wright, Felix Achana, Lavanya Diwakar, Malcolm G Semple, Will D Carroll, Kenneth Baillie, Christopher Thompson, Alice Alcock, Timothy S Kemp.

**Affiliations:**

University Hospitals of North Midlands NHS Trust, Stoke-on-Trent, UK (J. Wright, L. Diwakar, W.D Carroll, C. Thompson, A. Alcock, T.S Kemp).

University of Oxford, Oxford, UK (F. Achana).

University of Liverpool, Liverpool, UK (M.G Semple).

University of Edinburgh Roslin Institute, Edinburgh, UK (K. Baillie).

**Address for correspondence:**

Dr Jenny Wright, Department of Acute Internal Medicine, University Hospitals of North Midlands NHS Trust, Newcastle Road, Stoke-on-Trent, ST4 6QG, UK.

Email: [Jenny.Wright@uhn.nhs.net](mailto:Jenny.Wright@uhn.nhs.net)

Telephone: 01782715444

**Keywords:**

Covid-19, Respiratory Infection, Viral Infection

**Highlights**

- Multivariable regression analysis of data for hospitalised COVID-19 patients.
- We studied the likelihood of death and admission Ct values.
- Known clinical risk factors for the disease were adjusted for.
- Lower Ct values were associated with poorer outcomes in hospitalised patients.

**Unstructured abstract:**

In this single centre observational study, we demonstrated that lower cycle threshold (Ct) values (indicating higher viral loads) on admission to hospital, were associated with poorer outcomes in unvaccinated, hospitalised patients with Covid-19. We prospectively collected demographic and outcome data on all adult patients who tested positively for SARS-CoV-2 on admission to the University Hospitals North Midlands (UHNM) NHS Trust between 1<sup>st</sup> February and 1<sup>st</sup> July 2020. Nasopharyngeal swab samples were obtained, and a valid Ct value determined for all patients using the Public Health England (PHE) validated Viasure® reverse transcription PCR assay on admission to hospital. Multivariable logistic regression results based on data from 618 individuals demonstrated a statistically significant inverse relationship between the odds of death and Ct values (adjusted odds ratio (aOR) 0.95, 95% CI 0.92 to 0.98, p-value 0.001). The association remained highly statistically significant after adjusting for known clinical risk factors for the disease.

**Word count:**

852 words (excluding title, abstract, table/figures and references).

Clinicians need to identify patients with a higher risk of poor outcome and mortality from Covid-19 at an early stage of hospital admission. We describe a prospective, observational cohort study conducted at a UK tertiary care hospital to determine the relationship between the likelihood of death and Ct values in an unvaccinated UK population with Covid-19 on admission to hospital. Statistical adjustment was made for other known risk factors associated with poor outcome. Cycle thresholds (Ct) are semi-quantitative values, which are inversely proportional to the viral load in a reverse transcription polymerase chain reaction (PCR) test for SARS-CoV-2.

We prospectively collected demographic and outcome data on all adult patients who tested positively for SARS-CoV-2 on admission to the University Hospitals North Midlands (UHNM) NHS Trust between 1<sup>st</sup> February and 1<sup>st</sup> July 2020 using the ISARIC WHO Clinical Characterisation Protocol (Docherty et al., 2020). Nasopharyngeal swab samples were obtained, and a valid Ct value determined for all patients using the Public Health England (PHE) validated Viasure© reverse transcription PCR assay on admission to hospital. Ct

values below 38 were considered positive. Viasure© targets the *ORF1ab* (Target 1) and *N* genes (Target 2) of the SARS-CoV-2 viral genome (Public Health England, 2020). Target 1 was chosen as the primary exposure of interest as this is specific to SARS-CoV-2 viral genome. We combined Ct values with ISARIC data for statistical analysis.

Multivariable logistic regression was used to determine the association between Ct value and death within 28 days of a positive SARS-CoV-2 PCR test. We adjusted for age, gender, ethnicity, obesity and the presence of cardiovascular disease, chronic pulmonary disease, chronic kidney disease and diabetes. These covariates were selected apriori based on current understanding of the factors predicting poor outcomes in hospital cases of Covid-19 (Knight et al., 2020). Statistical significance was assessed at the 5% significance level and results presented as odds ratios and 95% confidence intervals. Analyses were conducted in R version 4.0.1.

A total of 803 SARS-CoV-2 positive adults (>18 years if age) with a valid Ct value determined on admission to hospital were eligible for inclusion in this study and were followed up for a period of up to 28 days [Figure 1]. The median age was 77 years (age range 19 to 100), 55% were male and 91% were of a white ethnicity. 35% had a history of cardiovascular disease, 18% chronic pulmonary disease, 19% had chronic kidney disease, 14% were asthmatic, 13% had either complicated and uncomplicated diabetes, and 7% were obese. The median cycle threshold value for target 1 was 25.8 (interquartile range 21.3 to 30.0). 285/803 patients (35.5%) died within 28 days of a positive SARS-CoV-2 PCR result.

Multivariable logistic regression results based on data from 618 (77.0%) individuals with complete covariate information are presented in Table 1. There was a statistically significant inverse relationship between the probability of death and Ct values (adjusted odds ratio [aOR 0.95(95% CI 0.92 to 0.98), p-value 0.001]. The only other variables that were significantly associated with mortality were age [aOR 1.05 (95% CI: 1.03 to 1.06), p-value <0.001] and diabetes [aOR 1.73 (95%CI: 1.01 to 2.97), p-value 0.044]


Sensitivity analyses carried after imputing missing data produced results consistent with the complete case analysis for both outcomes. Similar results were obtained when target 2 was used in the analysis instead of target 1 (these data are available on request).

In this prospective, observational study carried out at a tertiary UK hospital, lower Ct values (indicating higher viral loads) were associated with poorer outcomes in hospitalised patients with SARS-CoV-2 infection. This association remains highly statistically significant after adjusting for known clinical risk factors for the disease. High viral loads are associated with adverse outcomes in HIV and Ebola infections (Fitzpatrick et al., 2015; Li et al., 2015). There have been reports that higher viral loads are associated with adverse outcomes in Covid-19 patients in China, Taiwan and Brazil (Choudhuri et al., 2020; Faíco-Filho et al., 2020; Huang et al., 2020; Liu et al., 2020; Yu et al., 2020) but none from the UK.

The main strength of this study is that it uses a validated, generalisable dataset for analysis. Ct values were measured by validated methods. Robust statistical methods were used to account for interactions and missing data. Our study had several limitations. It was limited by its relatively small size. We therefore restricted our analysis to eight covariates known to be associated with poorer outcomes. Whilst this reduces the risk of confounding due to multiple  
prospective, observational cohort study conducted at a UK tertiary hospital.

hypothesis testing it did not allow exploration of the data to determine whether other measurements, such as admission cardiovascular status, oxygen saturation and liver failure or need for mechanical ventilation were associated with death. However, a logistic regression model incorporating all of these did not show any statistically significant associations. Moreover, the data is limited to a single tertiary care centre in the UK and the population characteristics may vary in relation to other areas. It is also unclear if these associations will remain significant after accounting for therapeutic agents (e.g., dexamethasone, anti-IL-6 treatment) in Covid-19 management. Our findings suggest that Ct values may have the potential to help clinicians identify patients at high risk of mortality from COVID-19 at the point of admission to hospital. We suggest that the analysis be repeated using a larger, multicentre dataset across different time periods.





versely associated with poorer outcomes in hospitalised patients with Covid-19: a short study conducted at a UK tertiary hospital.

Figure 1: A flow diagram illustrating how the final study sample size was determined.

Variable	Death		Death or continuous hospitalisation	
	Odds ratio (95% CI)	P-value	Odds ratio (95% CI)	P-value
Age (years)	1.05 (1.03, 1.06)	<0.001	1.04 (1.02, 1.05)	<0.001
Male	1.43 (1.00, 2.07)	0.052	1.2 (0.85, 1.69)	0.308
Non-White	0.66 (0.15, 2.21)	0.539	0.77 (0.23, 2.27)	0.65
Cardiovascular disease	1.00 (0.68, 1.47)	0.992	0.95 (0.65, 1.39)	0.799

Chronic pulmonary disease	1.17 (0.75, 1.82)	0.475	1.17 (0.76, 1.82)	0.469
Chronic kidney disease	1.22 (0.78, 1.91)	0.382	1.51 (0.97, 2.36)	0.071
Diabetes	1.68 (1.03, 2.75)	0.037	1.35 (0.83, 2.22)	0.227
Obesity	1.33 (0.68, 2.53)	0.39	1.1 (0.59, 2.05)	0.764
Cycle threshold	0.95 (0.92, 0.98)	0.001	0.94 (0.91, 0.97)	<0.001
Constant	0.01 (0.00, 0.04)	<0.001	0.05 (0.02, 0.12)	<0.001

Table 1: Multivariable logistic regression results based on complete case analysis

**Contributor statement:**

TK and JW conceived this study and analysis. MGS, JKB and ISARIC4C investigators [<https://isaric4c.net/about/authors/>] conceived the ISARIC WHO CCP-UK study. TK, JW, LD, AA, WC and CT contributed to the study design. TK and JW obtained ethical approval for this study and analysis. AA, JW and ISARIC4C investigators collected data for the study. FA conducted the primary statistical analysis. All authors contributed to writing and editing the final manuscript.

**HRA Ethical Approval:**

Health Research Authority (HRA) approval for this study and analysis was issued by the London - Chelsea Research Ethics Committee (REC reference: 20/HRA/3967) on 10<sup>th</sup> November 2020. Ethical approval for data collection by ISARIC4C in England was given by the South Central-Oxford C Research Ethics Committee in England (reference 13/SC/0149). The ISARIC WHO CCP-UK study was registered at <https://www.isrctn.com/ISRCTN66726260> and designated an Urgent Public Health Research Study by NIHR.

**Sponsor Organisation:**

University Hospitals of North Midlands NHS Trust

**Clinicaltrials.gov Identifier:**

Cycle threshold values are inversely associated with poorer outcomes in hospitalised patients with Covid-19: a prospective, observational cohort study conducted at a UK tertiary hospital.

NCT04651413

### **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### **Funding:**

No external funding was sought for this analysis. ISARIC4C is supported by grants from: The National Institute for Health Research (NIHR) [award CO-CIN-01], the Medical Research Council [grant MC\_PC\_19059].

### **Acknowledgments:**

This work uses data provided by patients and collected by the NHS as part of their care and support #DataSavesLives

## References

Choudhuri J, Carter J, Nelson R, Skalina K, Osterbur-Badhey M, Johnston A, Goldstein D, Paroder M, Szymanski J. SARS-CoV-2 PCR cycle threshold at hospital admission associated with patient mortality. PLoS One. 2020 Dec 31;15(12):e0244777. doi: 10.1371/journal.pone.0244777.

Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L, Holden KA, Read JM, Dondelinger F, Carson G, Merson L, Lee J, Plotkin D, Sigfrid L, Halpin S, Jackson C, Gamble C, Horby PW, Nguyen-Van-Tam JS, Ho A, Russell CD, Dunning J, Openshaw PJ, Baillie JK, Semple MG; ISARIC4C investigators. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation

Protocol: prospective observational cohort study. *BMJ*. 2020 May 22;369:m1985. doi: 10.1136/bmj.m1985.

Faíco-Filho KS, Passarelli VC, Bellei N. Is Higher Viral Load in SARS-CoV-2 Associated with Death? *Am J Trop Med Hyg*. 2020 Nov;103(5):2019-2021. doi: 10.4269/ajtmh.20-0954.

Fitzpatrick G, Vogt F, Moi Gbabai OB, Decroo T, Keane M, De Clerck H, Grolla A, Brechard R, Stinson K, Van Herp M. The Contribution of Ebola Viral Load at Admission and Other Patient Characteristics to Mortality in a Médecins Sans Frontières Ebola Case Management Centre, Kailahun, Sierra Leone, June-October 2014. *J Infect Dis*. 2015 Dec 1;212(11):1752-8. doi: 10.1093/infdis/jiv304. Epub 2015 May 22.

Huang JT, Ran RX, Lv ZH, Feng LN, Ran CY, Tong YQ, Li D, Su HW, Zhu CL, Qiu SL, Yang J, Xiao MY, Liu MJ, Yang YT, Liu SM, Li Y. Chronological Changes of Viral Shedding in Adult Inpatients With COVID-19 in Wuhan, China. *Clin Infect Dis*. 2020 Nov 19;71(16):2158-2166. doi: 10.1093/cid/ciaa631.

Knight SR, Ho A, Pius R, Buchan I, Carson G, Drake TM, Dunning J, Fairfield CJ, Gamble C, Green CA, Gupta R, Halpin S, Hardwick HE, Holden KA, Horby PW, Jackson C, Mclean KA, Merson L, Nguyen-Van-Tam JS, Norman L, Noursadeghi M, Olliaro PL, Pritchard MG, Russell CD, Shaw CA, Sheikh A, Solomon T, Sudlow C, Swann OV, Turtle LC, Openshaw PJ, Baillie JK, Semple MG, Docherty AB, Harrison EM; ISARIC4C investigators. Risk stratification of patients admitted to hospital with covid-19 using the

ISARIC WHO Clinical Characterisation Protocol: development and validation of the 4C Mortality Score. *BMJ*. 2020 Sep 9;370:m3339. doi: 10.1136/bmj.m3339. Erratum in: *BMJ*. 2020 Nov 13;371:m4334.

Li J, Duan HJ, Chen HY, Ji YJ, Zhang X, Rong YH, Xu Z, Sun LJ, Zhang JY, Liu LM, Jin B, Zhang J, Du N, Su HB, Teng GJ, Yuan Y, Qin EQ, Jia HJ, Wang S, Guo TS, Wang Y, Mu JS, Yan T, Li ZW, Dong Z, Nie WM, Jiang TJ, Li C, Gao XD, Ji D, Zhuang YJ, Li L, Wang LF, Li WG, Duan XZ, Lu YY, Sun ZQ, Kanu ABJ, Koroma SM, Zhao M, Ji JS, Wang FS. Age and Ebola viral load correlate with mortality and survival time in 288 Ebola virus disease patients. *Int J Infect Dis*. 2016 Jan;42:34-39. doi: 10.1016/j.ijid.2015.10.021.

Liu Y, Yan LM, Wan L, Xiang TX, Le A, Liu JM, Peiris M, Poon LLM, Zhang W. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis*. 2020 Jun;20(6):656-657. doi: 10.1016/S1473-3099(20)30232-2. Epub 2020 Mar 19.

Public Health England, Rapid assessment of the (ProLab/Certest) ViaSure SARS-CoV-2 Real Time PCR detection Kit (VS-NCO296T) .

[https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/889340/Rapid\\_assessment\\_ViaSure\\_SARS-CoV-2\\_Real\\_Time\\_PCR\\_detection\\_Kit\\_VS-NCO296T.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/889340/Rapid_assessment_ViaSure_SARS-CoV-2_Real_Time_PCR_detection_Kit_VS-NCO296T.pdf), 2020 (accessed April 2021).



Yu X, Sun S, Shi Y, Wang H, Zhao R, Sheng J. SARS-CoV-2 viral load in sputum correlates with risk of COVID-19 progression. Crit Care. 2020 Apr 23;24(1):170. doi: 10.1186/s13054-020-02893-8.

Journal Pre-proof